

## FUNGAL GLUCURONOYL AND FERULOYL ESTERASES FOR WOOD PROCESSING AND PHENOLIC ACID ESTER/SUGAR ESTER SYNTHESIS

Silvia Hüttner<sup>a, \*</sup>, Sylvia Klaubauf<sup>b</sup>, Hampus Sunner<sup>b</sup>, Cyrielle Bonzom<sup>a</sup>, Peter Jütten<sup>c</sup>, and Lisbeth Olsson<sup>a, b</sup>

<sup>a</sup> Department of Biology and Biological Engineering, Division of Industrial Biotechnology, Chalmers University of Technology, SE-412 96 Gothenburg, Sweden

<sup>b</sup> Wallenberg Wood Science Centre, Department of Biology and Biological Engineering, Chalmers University of Technology, SE-412 96 Gothenburg, Sweden

<sup>c</sup> Taros Chemicals GmbH & Co. KG, 44227 Dortmund

Feruloyl esterases (FAEs, E.C. 3.1.1.73, CAZy family CE1) and glucuronoyl esterases (GEs, E.C. 3.1.1.-, CAZy family CE15) are involved in the degradation of plant biomass by hydrolysing ester linkages in plant cell walls, and thus have potential use in biofuel production from lignocellulosic materials and in biorefinery applications with the aim of developing new wood-based compounds [1, 2]. GEs and FAEs are present in the genomes of a wide range of fungi and bacteria.

Under conditions of low water content, these enzymes can also carry out (trans)esterification reactions, making them promising biocatalysts for the modification of compounds with applications in the food, cosmetic and pharmaceutical industry. Compared to the chemical process, enzymatic synthesis can be carried out under lower process temperatures (50-60°C) and results in fewer side products, thus reducing the environmental impact.

We characterised new FAE and GE enzymes from mesophilic, thermophilic and cold-tolerant filamentous fungi produced in *Pichia pastoris*. The enzymes were characterised for both their hydrolytic abilities on various model substrates (methyl ferulate, pNP-ferulate) - for potential applications in deconstruction of lignocellulosic materials and extraction of valuable compounds - as well as for their biosynthetic capacities. We tested and optimised the FAEs' transesterification capabilities on ferulate esters in a 1-butanol-buffer system, with the aim of using the most promising candidates for the production of antioxidant compounds with improved hydrophobic or hydrophilic properties, such as prenyl ferulate, prenyl caffeate, glyceryl ferulate and 5-O-(trans-feruloyl)-arabinofuranose.

*Acknowledgments: Esterase research at Chalmers is supported by the EU Framework 7 project OPTIBIOCAT, the Knut and Alice Wallenberg Foundation (Wallenberg Wood Science Center) and the Swedish Research Council (Linnaeus Centre for Bio-inspired Supramolecular Function and Design, SUPRA).*

- [1] Topakas, E., Vafiadi, C., and Christakopoulos, P. (2007). Microbial production, characterization and applications of feruloyl esterases. *Process Biochemistry*, 42(4), 497–509. doi:10.1016/j.procbio.2007.01.007
- [2] Spániková, S., and Biely, P. (2006). Glucuronoyl esterase - Novel carbohydrate esterase produced by *Schizophyllum commune*. *FEBS Letters*, 580(19), 4597–601. doi:10.1016/j.febslet.2006.07.033